

EXHIBIT B

2)

1

Repeat WT-1 peptide binding Assay.

5x10⁵ T2 cells/well in RPMI + 5% Boiled FCS.Add peptides to a final conc. of 100 μ M.

Leave to binding at 37°C x 1

wash x 1 with PBS + 2% FCS.

Apply mAb A2 mAb HB34. HB117 30' on ice

wash x 3

Apply GAM FITC (1/250)

30' on ice

wash x 3

Resuspend cells in PBS (400 μ l).

FACS Analysis.

Results, Good.

peptide

Mean: HB34

HB117

- mAb only

513

37.70

42.32

HPV E7/86-93

109.70

118.12

HBV

146.86

220.67

WT-1 / 10-

81.62

144.38

WT-1 / 17-

43.79

54.04

WT-1 / 126-

69.78

30.79

WT-1 / 187-

93.01

136.27

WT-1 / 225-

70.03

97.56

WT-1 / 235-

95.63

105.90

WT-1 / 286-

45.66

76.42

WT-1 / 441-

137.32

172.31

BEST AVAILABLE COPY

A2 Binding Titration

12

peptides

WT:

10

17

126

187

225

235

280

441

HPVE7 EG-93

HBV 17-28

Plate peptide in RPMI + 5% Bolel FCS in 96 well plate (U bottomed), from 100 μ M of first column to Column 11 0.0001 μ M) in 50 μ l medium.

Add 50 μ l of T2 cells at a conc of 6×10^6 / well.
 5×10^5 / well mix well.

37°C overnight.

Wash x 1 w PBC (cold) + 1% FCS

Stain w HB 117 and HB 54 (noant culture sp) 10 min.
 for 30' on ice.

Wash x 3


Apply GAM Ig FITC at 1/250 (10 μ l)
 for 30' on ice

Wash x 3


Analyse on FACS

Results: Good see attached sheets.

WT-1 peptides binding to A2 molecules on T2 cells
 ----- Stained with HB54 mAb



Binding of WT-1 peptides to A2 molecules on T2 cells
 ----- Stained with HB117 mAb



3.)

4

Setting up Allo- & Self restricted peptide CTL lines

APMC: LG3 Az⁻ Allo

peptides: LG4 Az⁺ Self

WT1 10, 235, 441 and C228 (+ve Control).

Method: as left use DC from LG4 as stimulator

Pro stimulate with peptide loaded CIRAZ cells (50 min).

Restimulated E

" RMA-S-A2 (control)

11

CTL assay

Expand & cloning

Set up clones

Transfer to a well to 96 well plate

Feed

15

Setting up Self & Allo - restricted peptide specific
CTL line.

PBMC L G 5 - A2⁺ Self
L G 6 - A2⁻ Allo

pepptides: WT-1 126, 187, 280
+ve control E786-95.

Methods, see previous page.

First round stimulation with T2 loaded peptides.

Restimulate with peptide loaded - CIR A2

4/2 - RMA-S - A2

5/2 - ~~Restimulation~~ - CIR A2

7/3 - CTL

28/3 - Cloning

10/4 - CTL of clones

16

Protocol for peptide specific allorestricted human CTL

1. Prepare stimulator cells: in case of T2 stimulators preincubate o/n 10^6 cells/ml/well in x24 well plates in RPMI medium plus 5% boiled (10 min) FCS with 100 nM peptide. If allo DC are used as stimulators, allo DC from A2 +ve individuals can be separated as following:

Incubate A2+ PBMC at 37°C O/N, harvest nonadherent cells, resuspend in RPMI+9%FCS at 5×10^6 /ml and layer onto metrizadime gradient carefully, spin at 1500 for 10 min and harvest the interface (30% low density DC, 20% recovered from PBMC counts). Wash X3 with medium, resuspend at 10^6 /ml, add peptides to a final conc of 50uM and incubate at 37°C for 13000 rad. The stimulator can be added to responders directly w/o washing off peptides.

2. Separate blood by Ficoll centrifugation, wash twice, count, make sure that cells are HLA-A2 negative (by staining with HB54 and HB117 antibodies)

3. Plate 2×10^6 effectors and 2×10^5 stimulators per well in x24 well plates in culture medium (RPMI 1640, 10% FCS, 1% glutamine, 1% pen/strept 5×10^{-5} M 2-ME) with 500 nM peptide

4. On day 4 prepare stimulators as described in 1. if required. It may be not necessary to preincubate C1RA2 cells for more than 2 hrs

5. On day 5 harvest cells, count spin and seed 5×10^5 effectors, 2×10^6 irradiated (about 3000 rad) HLA-A2 negative autologous feeders, 2×10^5 peptide loaded and irradiated (about 10000 rad) T2 or C1RA2 cells and 500 nM peptide in 2ml of the culture medium with 10u/ml IL-2, 2.5ng/ml IL7 (1/2000 of the stock of 5ug/ml) and with 10% of Q12054 culture sup (anti-CD4 antibodies) (complete medium)

6. 14 days (and further every 2 weeks) later restimulate cultures as described in 5. Cell recovery is usually 200%


7. On day 5 analyse peptide specificity in a ^{51}Cr release assay using T2 or mouse cells transfected with A2 molecule as CTL targets and cold K562 cells as NK targets (10 per 1 labelled target). Use 0.7% TFA for max release instead of 0.5% SDS. Significant peptide specificity is detectable after 5-7 week in culture with E:T ratio 10:1

8. Peptide specific cultures can be cloned seeding 1, 10 and 100 cells/well in round bottom x96 well plates in complete culture medium with 10^4


18

Sheet1


allo LG6 wt126 specific CTL 24 well culture




allo LG6 wt187 specific ctl 24 well culture



alloLG6 wt280 24w



AII LG6 E7/86 CTL 24 well



572

-9
-16

67

-19

test clones 31 C Release Assay
Standard 4 hr


target T2 + E.7
T2 + 126
K562
K562 Az

more tests


Sheet1

10


hWT126 clone77




hWT126 clone 81




HWT126 CLONE85




HWT126 CLONE62



HWT126 CLONE 65



HWT126 CLONE1



771	-3	1199.2	4
907.4	0	979.2	0
	-3		74
	-2		79
	-2		25
	0		91

7)

11

51 Cr assay

clones 77. 81. 32.

targets: T2 + E7 as control

T2 + 126

Br 173

MV 441

G97

CARL Bonet Marrow 4 (Ferguson)

CD34+ from BM (by Steve)

E = T 25. 12. 6. 3. 1.5 075

Results:

line

line 32.

77

+

-81

T2 + E7

+

+

+

T2 + 126

+

+

+

Br 173

+

+

+

G97

+

+

+

MV 441

+

+

+

CARL CD34+

+

+

-

CARL BM 4

-

+

-


~~CD34+~~

The clone 77 kills CD34+

from CARL BM

12

HWT12 clone 77 @CD34 CMLBM



			-1		41
			-4		12
			-1		10
			-1		23
			28		27
			57		76
697	RNA	PCR	19		69
BV173		+	50		29
Ferguson		++	-2		80
Thomas	CD34+	+	-2		0
Grady		+	-4		11
		-	-2		60
			-1		68
			67		40
			13		31
			6		97
			3		97
			0		100
			-5		105
			-5		19
			-7		117
			13		99
			-3		100
			-5		108
					25
					85
					102
					61
					88
					99
					87
					12
					61
					25
					101
					89
					24
					81
					85
					26
					118
					26
					28
					46
					96
					69
					1
					76

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.